

CAUTION

Harmful if swallowed. IF SWALLOWED: Call a POISON CENTER, doctor, physician or first aider if you feel unwell. Causes skin irritation. IF ON SKIN: Wash with plenty of water. If skin irritation occurs: Get medical advice. Causes serious eye damage. IF IN EYES: Rinse cautiously with water for several minutes. Harmful if inhaled. IF INHALED: Remove person to fresh air and keep comfortable for breathing. Wash all exposed external body areas thoroughly after handling.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone **Australia: 13 11 26; New Zealand 0800 764 766.**

**IN AN EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE**

**CROP SMART PTY LTD
EMERGENCY 1300 783 481
AUSTRALIA WIDE, 24 HOURS**

Scan QR Code to access SDS
on mobile device



CONDITIONS OF SALE

Crop Smart Pty Ltd ("Crop Smart") shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever, or howsoever arising through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express conditions that the purchaser does not rely on Crop Smart's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Crop Smart has any authority to alter these conditions.

Crop Smart Pty Ltd
ABN 28 093 927 961
2409/4 Daydream Street,
Warriewood NSW 2102
Tel: 1300 783 481 Fax: 1300 783 491

DANGER
KEEP OUT OF REACH OF CHILDREN

Mallee Mix Sulphate

Smart Sulphate

Manufactured using water soluble sulphates.
Product solutions are maximised with
elemental value.

Typical Analysis (W/V)%
Manganese (Mn).....6
Zinc (Zn) 4.8
Copper (Cu) 1.2
Sulphur (S)7
Molybdenum (Mo) 0.025

Crop Smart
better crop protection



APPLICATION RATES

Broadacre Crops	Tree Crop / Vegetables / Vines
Foliar 1-4L application rate per hectare	Foliar 1-4L application rate per hectare
Infurrow 3-8L application rate per hectare	Fertigation 3-8L application rate per hectare
Aerial Application Not recommended	Aerial Application Not recommended

Store out of direct sunlight and mix well prior to use.

Foliar fertilisers work best when applied in cool dry conditions. Do not apply as a foliar when temperatures exceed 25 degrees Celsius. Do not apply as a foliar fertiliser if crop is more than 20% flowering.

COMPATIBILITY

Smart Mallee Mix Sulphate is compatible with most agricultural chemicals and fertilisers. Do not mix with Amine herbicides and Calcium containing products. When mixing, always test for compatibility with a small quantity first and only mix enough for one application. Do not store the mixed product. Do not mix with inoculants and alkaline products.

WARNING

Excessive use of molybdenum (Mo) can be harmful to stock. Plant levels of Mo can be high for 4 weeks after application. It is advisable to keep stock off treated areas for this period. Molybdenum application may induce copper deficiency in grazing animals. This is most likely to occur on sandy soils low in copper. Check rate and frequency of molybdenum use with appropriate authorities.

Batch Number:

Date of Manufacture:

20L 110L 1000L