### **CAUTION**

a POISON CENTER, doctor, physician or first aider if you feel unwell. Causes skin irritation. Causes serious eye damage. IF IN EYES: Rinse cautiously with water for several minutes. May cause damage to organs through prolonged or repeated exposure.

# FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. Phone Australia: 13 11 26; New Zealand 0800 764 766.

> **IN AN EMERGENCY DIAL 000 POLICE OR FIRE BRIGADE**

**CROP SMART PTY LTD EMERGENCY 1300 783 481 AUSTRALIA WIDE, 24 HOURS** 

Scan QR Code to access SDS on mobile device



## **CONDITIONS OF SALE**


Crop Smart Ptv Ltd ("Crop Smart") shall not whether consequential or otherwise whatsoever, or howsoever arising through negligence or use or application of this product. The supply of this product is on the express conditions that the judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Crop Smart has any authority to alter these conditions.

Crop Smart Pty. Ltd

ABN 28 093 927 961 2409/4 Daydream Street Warriewood NSW 2102 Tel: 1300 783 481 Fax: 1300 783 491

# **DANGER KEEP OUT OF REACH OF CHILDREN**

# ZMC plus Moly Sulphate 6.1-5.1-1.5



# **APPLICATION RATES**

Broadacre Crops	Tree Crop / Vegetables / Vines
Foliar 1-4L application rate	Foliar 1-4L application rate
per hectare Infurrow	per hectare Fertigation
3-8L application rate per hectare	3-8L application rate per hectare
Aerial Application Not recommended	Aerial Application Not recommended

# Store out of direct sunlight and mix well prior to use.

Foliar fertilisers work best when applied in cool dry conditions. Do not apply as a foliar when temperatures exceed 25 degrees Celsius. Do not apply as a foliar fertiliser if crop is more than 20% flowering.

# **COMPATIBILITY**

Smart ZMC plus Moly Sulphate 6.1-5.1-1.5 is compatible with most agricultural chemicals and fertilisers. Do not mix with Amine herbicides and test for compatibility with a small quantity first and only mix enough for one application. Do not store the mixed product. Do not mix with inoculants and alkaline products.

### **WARNING**

Excessive use of molybdenum (Mo) can be harmful to stock. Plant levels of Mo can be high for 4 weeks off treated areas for this period. Molybdenum grazing animals. This is most likely to occur on sandy soils low in copper. Check rate and frequency of molybdenum use with appropriate authorities.

Batch Number:
Date of Manufacture:

■ 20L ■ 110L ■ 1000L



