

**POISON**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# Smart Azure 500

## FUNGICIDE

**ACTIVE CONSTITUENT: 500 g/L FLUTRIAFOL**

**GROUP 3 FUNGICIDE**

For control of certain fungal diseases on wheat, barley and canola when mixed with fertiliser as per the Directions for Use.

**IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USING THIS PRODUCT**

Batch Number:

Date of Manufacture:

Scan QR Code to  
access SDS  
on mobile device



APVMA Approval Number: 94139/141676

**Crop Smart**  
better crop protection

Crop Smart Pty. Ltd  
ABN 28 093 927 961  
2409/4 Daydream Street, Warriewood NSW 2102  
Tel: 1300 783 481 Fax: 1300 783 491



**POISON**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

# Smart Azure 500

## FUNGICIDE

**ACTIVE CONSTITUENT: 500 g/L FLUTRIAFOL**

**GROUP 3 FUNGICIDE**

For control of certain fungal diseases on wheat, barley and canola when mixed with fertiliser as per the Directions for Use.

**IMPORTANT: READ THIS LEAFLET BEFORE USE**

**CONTENTS: 10L – 1000L**

Crop Smart Pty. Ltd  
ABN 28 093 927 961  
2409/4 Daydream Street, Warriewood NSW 2102  
Tel: 1300 783 481 Fax: 1300 783 491

APVMA Approval Number: 94139/141676

**Crop Smart**  
better crop protection



## DIRECTIONS FOR USE: All States

Crop	Disease	Rate per hectare	WHP	Critical Comments																					
Canola	Blackleg ( <i>Leptosphaeria maculans</i> )	200 mL	4 weeks Grazing	Treatment will reduce blackleg infection in both susceptible and tolerant canola varieties. <b>Application:</b> Spray undiluted product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate, which will deposit 200 mL of the product on the quantity of fertiliser required to be sown in furrow per hectare, using the table below as a guide.																					
				<table border="0"> <thead> <tr> <th>Fertiliser rate per hectare (kg)</th> <th>Volume of product per 100 kg of fertiliser (mL)</th> </tr> </thead> <tbody> <tr> <td>60</td> <td>333*</td> </tr> <tr> <td>70</td> <td>285</td> </tr> <tr> <td>80</td> <td>250</td> </tr> <tr> <td>100</td> <td>200</td> </tr> <tr> <td>120</td> <td>166</td> </tr> </tbody> </table>	Fertiliser rate per hectare (kg)	Volume of product per 100 kg of fertiliser (mL)	60	333*	70	285	80	250	100	200	120	166									
Fertiliser rate per hectare (kg)	Volume of product per 100 kg of fertiliser (mL)																								
60	333*																								
70	285																								
80	250																								
100	200																								
120	166																								
Wheat	Stripe rust ( <i>Puccinia striiformis</i> )	100 or 200 mL		For control of stripe rust of wheat and control of powdery mildew and scald of barley, use the high rate where conditions conducive to severe disease are expected. <b>Application:</b> Spray product evenly onto granulated fertiliser and immediately mix thoroughly in a mechanical mixer or auger. Apply at a rate which will deposit the specific rate of the product on the quantity of fertiliser required to be sown in-furrow per hectare, using the table below as a guide.																					
	Septoria tritici blotch ( <i>S. tritici</i> ) Take-all ( <i>Geaumannomyces graminis</i> )	200 mL		<table border="0"> <thead> <tr> <th>Fertiliser rate per hectare (kg)</th> <th>Volume of product per 100 kg of fertiliser (mL)</th> </tr> </thead> <tbody> <tr> <td></td> <td>100 mL rate</td> <td>200 mL rate</td> </tr> <tr> <td>50</td> <td>200</td> <td>400 *</td> </tr> <tr> <td>60</td> <td>166</td> <td>333 *</td> </tr> <tr> <td>70</td> <td>143</td> <td>285</td> </tr> <tr> <td>80</td> <td>125 †</td> <td>250</td> </tr> <tr> <td>100</td> <td>100 †</td> <td>200</td> </tr> <tr> <td>120</td> <td>88 †</td> <td>166</td> </tr> </tbody> </table>	Fertiliser rate per hectare (kg)	Volume of product per 100 kg of fertiliser (mL)		100 mL rate	200 mL rate	50	200	400 *	60	166	333 *	70	143	285	80	125 †	250	100	100 †	200	120
Fertiliser rate per hectare (kg)	Volume of product per 100 kg of fertiliser (mL)																								
	100 mL rate	200 mL rate																							
50	200	400 *																							
60	166	333 *																							
70	143	285																							
80	125 †	250																							
100	100 †	200																							
120	88 †	166																							
Barley	Powdery mildew ( <i>Erysiphe graminis</i> ) Scald ( <i>Rhynchosporium secalis</i> )	100 or 200 mL		* Application volumes over 300 mL per 100kg of fertiliser may adversely affect the flow rate and handling characteristics of dusty/poorly granulated fertilisers and some high nitrate fertilisers. † Where application volume is low, product may be diluted with water to improve coverage.																					

**NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**WITHHOLDING PERIODS: APPLIED IN-FURROW AS A FERTILISER TREATMENT**

**CANOLA, WHEAT AND BARLEY - DO NOT GRAZE OR CUT FOR STOCK FOOD FOR 4 WEEKS AFTER SOWING**

## **GENERAL INSTRUCTIONS**

### **FERTILISER APPLICATION**

Granulated Fertiliser

To determine the correct volume of Smart Azure 500 Fungicide required to be applied, refer to critical Comments in the Directions for Use table, which provides a guide based on a range of common fertiliser rates.

Ensure that treated fertiliser is dry before sowing with the crop.

Treated fertiliser should be applied in a band 3-4 cm below or to the side of the seed in the planting furrow. It should not be broadcast.

Note that the flow rate of fertiliser may be changed after treatment, and the seeder should be calibrated with the treated material to ensure that the correct rates of fertiliser and fungicide are delivered.

Do not mix seed with treated fertiliser and Smart Azure 500 Fungicide.

### **FUNGICIDE RESISTANCE WARNING**

#### **GROUP 3 FUNGICIDE**

Smart Azure 500 Fungicide is a member of the DMI group of fungicides. For fungicide resistance management Smart Azure 500 Fungicide is a Group 3 fungicide.

Some naturally occurring individual fungi resistant to Smart Azure 500 Fungicide and other Group 3 fungicides may exist through normal genetic variability in any fungal population.

The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Smart Azure 500 Fungicide or other Group 3 fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, Crop Smart Pty Ltd accepts no liability for any losses that may result from the failure of Smart Azure 500 Fungicide to control resistant fungi. Resistance should not be assumed without first reviewing the method of application, the coverage and the timing of application.

### **PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

DO NOT contaminate streams, rivers or watercourses with the chemical or used containers

### **PRECAUTIONS**

Re-entry or Re-handling

DO NOT allow entry into treated areas until the spray has dried, unless wearing cotton overalls buttoned to the neck and wrist (or equivalent clothing) and chemical resistant gloves. Clothing must be laundered after each day's use.

### **STORAGE AND DISPOSAL:**

Store in the closed, original container in a cool, well-ventilated area. Do not store for prolonged periods in direct sunlight.

Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site.

If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up

for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

For refillable containers: Empty contents fully into application equipment. Close all valves and return to point of supply for refill or storage.

#### **SAFETY DIRECTIONS**

Harmful if swallowed. Harmful if absorbed by skin contact. May irritate the skin. Will irritate the eyes. Avoid contact with eyes and skin. When opening the container, preparing product for use and using the product, wear cotton overalls buttoned to the neck and wrist, elbow-length chemical resistant gloves and face shield or goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves and face shield or goggles and contaminated clothing.

#### **FIRST AID INSTRUCTIONS:**

If poisoning occurs contact a doctor or Poisons Information Centre. Tel **Australia 13 11 26, New Zealand 0800 764 766**.

#### **SAFETY DATA SHEET**

For further information refer to the Safety Data Sheet (SDS).

#### **CONDITIONS OF SALE:**

Crop Smart Pty Ltd shall not be liable for any loss, injury, damage or death, whether consequential or otherwise whatsoever or howsoever arising whether through negligence or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Crop Smart's skill or judgement in purchasing or using the same and every person dealing with this product does so at his own risk absolutely. No representative of Crop Smart has any authority to add to or alter these conditions.



#### **ADDITIONAL GHS STATEMENT**

Harmful if swallowed. IF SWALLOWED: Call a POISON CENTER, doctor, physician or first aider if you feel unwell.

Wash all exposed external body areas thoroughly after handling. Do not eat, drink or smoke when using this product.

Harmful to aquatic life with long lasting effects. Avoid release to the environment.

**IN AN EMERGENCY  
DIAL 000  
POLICE OR FIRE BRIGADE**

**EMERGENCY CONTACT CHEMWATCH  
1800 951 288  
AUSTRALIA WIDE, 24 HOURS**

