

POISON

KEEP OUT OF REACH OF CHILDREN

READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Smart Team T

SEED DRESSING

ACTIVE CONSTITUENTS: 360 g/L THIRAM
200 g/L THIABENDAZOLE

GROUP **M3** | **1** FUNGICIDE

Controls seed-borne & seedling root pathogens of peas, chickpeas, lentils, faba beans and vetch as specified in the Directions for Use table.

IMPORTANT: READ THE ATTACHED BOOKLET BEFORE USING THIS PRODUCT

Crop Smart Pty. Ltd
ABN 28 093 927 961
2409/4 Daydream Street, Warriewood NSW 2102
Tel: 1300 783 481 Fax: 1300 783 491

Batch Number:

Date of Manufacture:

Scan QR Code to
access SDS
on mobile device



APVMA Approval Number: 88390/121187

Crop Smart

better crop protection



POISON
KEEP OUT OF REACH OF CHILDREN
READ SAFETY DIRECTIONS BEFORE OPENING OR USING

Smart
Team T

SEED DRESSING

**ACTIVE CONSTITUENTS: 360 g/L
THIRAM 200 g/L THIABENDAZOLE**

GROUP M3 1 FUNGICIDE

**Controls seed-borne & seedling root pathogens of peas,
chickpeas, lentils, faba beans and vetch as specified in the
Directions for Use table.**

IMPORTANT: READ THIS LEAFLET BEFORE USE

CONTENTS: 10L – 200L

Crop Smart Pty. Ltd
ABN 28 093 927 961
2409/4 Daydream Street, Warriewood NSW 2102
Tel: 1300 783 481 Fax: 1300 783 491



APVMA Approval Number: 88390/121187

Crop Smart
better crop protection

DIRECTIONS FOR USE (ALL STATES)

Restrains:

DO NOT mix this product with legume Rhizobium inoculant.

SEED TYPE	DISEASE	RATE	CRITICAL COMMENTS
Peas Field and Garden	Black Spot (Leaf and Pod Spot and Collar Rot) (<i>Mycosphaerella pinodes</i> , <i>Ascochyta pisi</i> , <i>Phoma medicaginis</i> , var. <i>pinodella</i>)	200mL per 100 kg of seed	<p>Peas, lentils, faba beans, vetch: This liquid seed dressing may be diluted with 1 part water to parts product. Apply 300 mL of diluted mixture to 100 kg of seed.</p> <p>Chickpeas: In the case of chickpea seed, mix 8 parts of water to 2 parts of product. Apply 1 L of diluted mixture to 100kg of chickpea seed.</p> <p>Seed must be treated uniformly using corrected calibrated seed treatment equipment. Secondary mixing of treated seed (e.g. in an auger) is important to obtain even coverage.</p> <p>In the case of <i>Ascochyta</i> (peas, chickpeas and lentils) and <i>Botrytis</i>(chickpeas), Smart Team T Seed Dressing will provide control of seed borne infection but follow up foliar fungicide sprays registered for this purpose, will be required to control field infection from other sources.</p>
	Seedling Root Rots (<i>Fusarium</i> spp., <i>Pythium</i> spp., * <i>Macrophomina phaseolina</i>)		
Chickpeas	<p><i>Ascochyta</i> Blight (<i>Ascochyta rabiei</i>), <i>Botrytis</i> Seed Rot (<i>Botrytis cinerea</i>), Seedling Root Rots (<i>Fusarium</i> spp., <i>Pythium</i> spp.)</p>		
Lentils	<p><i>Ascochyta</i> Blight (<i>Ascochyta lentis</i>), Seedling Root Rots (<i>Fusarium</i> spp., <i>Pythium</i> spp.)</p>		
Faba Beans, Vetch	Seedling Root Rots (<i>Fusarium</i> spp., <i>Pythium</i> spp.)		

*This disease is not known to occur in WA.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED.

GENERAL INSTRUCTIONS

This product does not adversely affect germination, however when Rhizobium cultures (inoculum) are also added to seed, the viability of the inoculum may be reduced. Always apply the fungicide first and allow to dry. Apply the inoculum immediately prior to sowing. Seed treatment will control Black Spot and Fusarium spp., Ascochyta spp. and Botrytis cinerea pathogens already present in pea seed.

Treatment will also give young pea plants early protection from any external infection from Black Spot.

Seed treatment can alter seed flow characteristics. The drill should be recalibrated using treated seed prior to seeding.

Equipment used for applying this product can be cleaned out with water. All seed that has been stored may lose viability especially if it becomes hot or damp. Smart Team T Seed Dressing has no adverse effect on seed viability or seedling vigour for up to 15 months after treatment, but all seed should be tested for germination in the year it is used to ensure that there is sufficient viable seed sown to establish the optimum seedling density.

Smart Team T Seed Dressing treated seed carried over from previous seasons may not provide adequate control of soil borne pathogens (eg. Pythium spp. and Fusarium spp.). Treated seed is to be used only for sowing.

MIXING

Shake well before use. Add water and product to the applicator tank and mix well. Maintain agitation during use. Where the diluted product has been permitted to stand for an extended period, the mixture should be again agitated well. DO NOT permit this product to dry-out in application equipment as flaking may occur and contribute to nozzle blockages.

INSECTICIDE RESISTANCE WARNING

GROUP	M3	1	FUNGICIDE
-------	-----------	----------	-----------

Smart Team T Seed Dressing is a combination of a Benzimidazole and a Multi-site Activity fungicide. For fungicide resistance management Smart Team T Seed Dressing is both a Group M3 and a Group 1 fungicide. Some naturally occurring individual fungi resistant to Smart Team T Seed Dressing and other Group M3 and/or Group 1 fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungi population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Smart Team T Seed Dressing and other Group M3 and/or Group 1 fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, Crop Smart Pty Ltd accepts no liability for any losses that may result from the failure of Smart Team T Seed Dressing to control resistant fungi.

PROTECTION OF LIVESTOCK

DO NOT allow treated seed to contaminate grain intended for animal consumption. Treated seed is not to be used as food for poultry or any other animal.

PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT

DO NOT feed treated seed or otherwise expose to wild or domestic birds. DO NOT contaminate dams, rivers, streams, ponds, waterways and drains with. Smart Team T Seed Dressing, with the used containers, or bags or other containers which have held treated seed.

STORAGE AND DISPOSAL

Store in the closed, original container in a cool, well-ventilated area. DO NOT store for prolonged periods in direct sunlight. Store in a locked room or place away from children, animals, food, feedstuffs, seed and fertilisers. DO NOT repack or use container for other purposes. When treated seed is stored it must be stored apart from other seed and bags or containers must be clearly marked indicating the contents have been treated with Smart Team T Seed Dressing. Treated seed is not to be used for human consumption. Bags or other containers which have held treated seed are not to be used for any other purpose.

For Non-Refillable Containers: Triple-rinse containers before disposal. Add rinsings to spray tank. Do not dispose of undiluted chemicals on site.

If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush, or puncture and deliver empty packaging to an approved waste management facility. If an approved waste management facility is not available, bury the empty packaging 500 mm below the surface in a disposal pit specifically marked and set up for this purpose, clear of waterways, desirable vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

For Refillable Containers: Storage must be secure so that contents cannot be tampered with. All locks and/or seals must be in order. If locks or seals are broken prior to initial use, then the integrity of this product cannot be assured. If this occurs Crop Smart Pty Ltd should be advised immediately. Empty contents into application equipment, close all valves and return to the point of supply for refill or storage.

SAFETY DIRECTIONS

May irritate the eyes, nose, throat and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. Do not inhale spray mist. When opening the container, preparing and using the product wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length PVC gloves and face shield or goggles. If product on skin, immediately wash area with soap and water. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs contact a doctor or Poisons Information Centre. **Phone Australia 13 11 26; New Zealand 0800 764 766.** Avoid giving alcohol.

SAFETY DATA SHEET

For further information refer to the Safety Data Sheet which is available from the supplier.

NOTICE TO BUYERS

Crop Smart Pty Ltd ('Crop Smart') shall not be liable for any loss, injury, damage or death whether consequential or otherwise whatsoever or howsoever arising whether through negligence, use under abnormal conditions or otherwise in connection with the sale, supply, use or application of this product. The supply of this product is on the express condition that the purchaser does not rely on Crop Smart's skill or judgment in purchasing or using the product and every person dealing with this product does so at their own risk.



ADDITIONAL GHS STATEMENT

Harmful if swallowed. IF SWALLOWED: Call a POISON CENTRE or doctor if you feel unwell. Causes skin irritation. IF ON SKIN: Wash with plenty of water. May cause respiratory irritation. IF exposed or concerned: Get medical advice or attention. Very toxic to aquatic life with long lasting effects. Avoid release to the environment.

**IN AN EMERGENCY
DIAL 000
POLICE OR FIRE BRIGADE**

**EMERGENCY CONTACT CHEMWATCH
1800 951 288
AUSTRALIA WIDE, 24 HOURS**

